

Carbon Monoxide Poisoning: Pathophysiology, Diagnosis, and Forensic Challenges in Fatal Exposures

Ariba Memon¹, Dr. Priti Patel¹

¹Bachelor of Pharmacy, Dr. L. H. Hiranandani College of Pharmacy, Thane, Maharashtra, India.

Corresponding author: priti.patel@dlhcop.org

Doi: <https://doi.org/10.5281/zenodo.22301925>

Received: 07th July 2026

Accepted: 17th July 2026

Abstract.

Carbon monoxide (CO) poisoning is a serious worldwide public health concern and the primary cause of unintentional toxic fatalities [5, 8, 15]. Many instances remain undiagnosed, leading to increased morbidity and death. As it is colorless, odorless, and non-irritating, it is also known as the “silent killer.” [5, 11] Its toxicity is largely due to carboxyhemoglobin production, which lowers oxygen supply, combined with mitochondrial dysfunction, oxidative stress, and inflammation, notably damaging the brain and heart [5, 8, 10]. The clinical presentation varies from modest symptoms such as headache and dizziness to severe neurological impairment, cardiac problems, coma, and death. [8, 11, 15] Diagnosis might be challenging since signs and symptoms are non-specific and the usual methods of COHb testing and imaging are limited, especially in fatal cases [1, 15]. This review covers CO toxicokinetics, etiology, clinical characteristics, diagnostics, and treatment options, including hyperbaric oxygen therapy [12, 14, 15]. It also discusses the delayed neurological sequelae and forensic aspects of CO poisoning, including autopsy results and medico-legal review [1, 2, 3], as well as novel biomarkers and preventative methods for future clinical and forensic practice.

Keywords: carbon monoxide poisoning, carboxyhemoglobin, hyperbaric oxygen treatment, forensic toxicology, delayed neurological sequelae, and biomarkers.

Introduction:

Carbon monoxide (CO) poisoning is a leading cause of preventable inhalational illness and death worldwide [5, 8, 15]. Exposure is often encountered in confined or inadequately ventilated areas owing to residential heating systems, engines, fires, and industrial sources [3, 4, 11]. People are generally unaware of exposure to CO until they have serious symptoms, since CO is colorless, odorless, and non-irritating; hence, it has a significant risk of death [5, 11, 8, 15]. CO toxicity has been related for a long time to the development of carboxyhemoglobin and decreased delivery of oxygen, but it also includes oxidative stress, mitochondrial dysfunction, inflammation, and damage to lipid membranes [5, 10, 8, 15]. Mechanisms of action are especially important in high oxygen demand organs such as the brain and heart and may cause delayed neurological consequences even after apparent recovery [5, 10, 7, 12]. The first symptoms, including headache, dizziness, exhaustion, and nausea, are vague and may resemble other illnesses, which can delay detection and complicate diagnosis [8, 11, 15]. In fatal instances, interpretation is further confounded by limits of postmortem carboxyhemoglobin assays and varied results [1, 2, 3]. This review provides an overview of the epidemiology, pathophysiology, diagnosis, treatment, and forensic aspects of CO poisoning with a focus on diagnostic problems, delayed neurological sequelae, and new therapeutic and investigative breakthroughs that can improve future outcomes [1, 12, 13, 15].

Epidemiology and Sources of Exposure:

Carbon monoxide (CO) poisoning is one of the primary causes of inhalation-related illness and death globally [3, 5]. It is commonly underdiagnosed or underreported in a number of nations; therefore, the real impact is larger than acknowledged [3, 4]. Cases are higher in winter months owing to increased indoor usage of combustion-based heating systems [3, 11]. Unintentional home exposures are the cause of most instances [3, 5].

Exposure may be purposeful or unintentional. Common causes include car exhausts, malfunctioning residential appliances, charcoal fires, generators, industrial combustion systems, and smoke from burning buildings, which may also include additional dangerous gases, including hydrogen cyanide [4, 5, 11].

The high-risk populations are youngsters, the elderly, pregnant women, and those with cardiovascular or respiratory problems [5, 8, 13]. Poor ventilation, malfunctioning equipment, absence of CO alarms, and low knowledge (particularly in under-resourced settings) significantly enhance risk [3, 4, 15]. In general, CO poisoning is a major environmental and public health problem [4, 15].

Table 1. Common Sources of Carbon Monoxide Exposure.

Category	Common Sources	Typical Setting
Domestic	Gas heaters, stoves, charcoal burners, generators	Homes and residential buildings
Occupational	Industrial furnaces, mining operations, vehicle repair workshops	Workplaces and industrial settings
Fire-related	Structural fires, smoke inhalation	Residential and commercial fires
Transportation	Vehicle exhaust emissions	Garages, tunnels, enclosed parking areas
Environmental	Poorly ventilated combustion sources	Confined or enclosed spaces

Toxicokinetics and Pathophysiology:

Carbon monoxide (CO) is readily absorbed in the lungs and enters the bloodstream, where it binds to hemoglobin 200 to 250 times more tightly than oxygen, creating carboxyhemoglobin (COHb) [5, 8]. This results in a decreased oxygen supply and a leftward shift of the oxyhemoglobin dissociation curve [5, 10]. The half-life of COHb depends on oxygen concentration and is greatly shortened by supplementary oxygen treatment [12, 14]. CO toxicity is not only hypoxia but also includes myoglobin, cytochromes, and other heme proteins. It disrupts mitochondrial oxidative phosphorylation via inhibition of cytochrome c oxidase [5, 10]. This leads to decreased cellular respiration, oxidative stress, and energy failure, which are important contributors to damage [8, 10, 15]. Inflammation also contributes to harm via leukocyte activation, endothelial dysfunction, and production of pro-inflammatory mediators and contributes to microvascular and neuronal injury and delayed neuropsychiatric sequelae [7,10,15]. The brain and heart are especially susceptible, with frequent observations of basal ganglia, hippocampus, and white matter injury, as well as myocardial ischemia, arrhythmias, and cardiac failure [7, 10, 13, 15].




Figure 1: Pathophysiologic Mechanisms of Carbon Monoxide Poisoning

Clinical Manifestations:

The symptoms of carbon monoxide (CO) poisoning depend on exposure and time and the individual susceptibility [8, 11, 15]. Initial symptoms are vague and include headache, dizziness, tiredness, nausea, vomiting, and muscular weakness, which commonly results in delayed diagnosis [5, 8]. As exposure increases, neurological symptoms such as disorientation, poor judgement, blurred vision, and lack of coordination may develop [8, 15]. In extreme situations, CO poisoning may induce convulsions, loss of consciousness, and coma [5, 8]. Cardiovascular consequences include tachycardia, chest discomfort, arrhythmias, myocardial ischemia, and general cardiac dysfunction [13, 15]. Respiratory symptoms may also be present, especially with smoke inhalation [5, 11]. High-risk people include pregnant women, children, the elderly, and those with cardiovascular disease [5, 8]. Importantly, the levels of COHb do not necessarily correspond with the severity of symptoms, and some patients have delayed neurocognitive consequences such as memory loss and mood abnormalities even after apparent recovery [7, 12, 15].

Table 2: Clinical Manifestations of Carbon Monoxide Poisoning

Severity	Common Manifestations
Mild	Headache, dizziness, fatigue, nausea, vomiting
Moderate	Weakness, confusion, impaired concentration, visual disturbances
Severe	Altered consciousness, seizures, coma, respiratory failure
Cardiovascular Effects	Chest pain, arrhythmias, myocardial ischemia, myocardial injury
Delayed Effects	Memory impairment, personality changes, cognitive dysfunction, movement disorders

Diagnosis and Biomarkers:

Diagnosis of CO poisoning is based on the history, physical examination, and laboratory work-up of the patient. COHb values are the gold standard for patient diagnosis. However, there are substantial challenges in interpretation. The COHb level may be reduced even before the visit to the physician, particularly when the patient was treated for the disease; smokers have elevated baseline COHb levels [5, 11, 15].

Lab tests may show metabolic acidosis, elevated lactate in blood, and evidence of damage to organs. For instance, cardiac markers such as troponins and creatine kinase may be helpful in evaluating myocardial involvement in CO poisoning and may be indicative of a bad prognosis in this regard [13, 15]. Furthermore, currently, neuroimaging tests using magnetic resonance imaging are usually conducted for the diagnosis of brain damage in severe poisoning [7].

The latest studies related to CO biomarkers include metabolomics and oxidative stress indices. These approaches are useful to forecast the prognosis of patients and to encourage the early diagnosis of severe CO poisoning [1, 15]. In situations of death, diagnosis should be made by thorough interpretation of postmortem COHb level, toxicological tests, and probable evidence of tissue damage by decomposition or injuries incurred during fire or co-toxicant exposure [1, 2, 3].

Table 3: Diagnostic Modalities and Biomarkers in Carbon Monoxide Poisoning

Diagnostic Tool / Biomarker	Clinical Significance	Limitation
Carboxyhemoglobin (COHb)	Confirms exposure to carbon monoxide	Levels may decrease before testing.
Arterial Blood Gas Analysis	Assesses oxygenation and acid-base status	Findings may be nonspecific
Serum Lactate	Indicates tissue hypoxia and poisoning severity	Not specific to CO poisoning
Cardiac Troponin	Detects myocardial injury and prognostic risk	Elevated in other cardiac conditions
Magnetic Resonance Imaging (MRI)	Identifies neurological injury and predicts sequelae	Expensive and not always available
Metabolomic Biomarkers	Emerging tool for prognosis and forensic evaluation	Requires further validation

Management and Treatment:

And so the critical thing to do when someone gets carbon monoxide poisoning is to get them away from the object that is creating the carbon monoxide immediately. Then we have to give them enough oxygen to breathe. This helps clear the carbon monoxide from their blood and ensures their body receives the oxygen it needs. [5, 12, 14].

Supportive treatment is especially important in moderate to severe instances. Management involves ensuring that the airway is clear, giving oxygen, and careful monitoring of cardiovascular and respiratory function to avoid complications. [5, 8, 14].

Hyperbaric oxygen therapy (HBOT) is a common treatment researched for carbon monoxide intoxication. HBOT provides 100% oxygen at increased ambient pressure and enhances carbon monoxide removal and tissue oxygenation [12, 14]. Potential advantages include reduced neuronal injury, limitation of inflammatory damage, and prevention of later neurological sequelae. However, the clinical usage has been variable, with ambiguity about patient selection, timing, and long-term efficacy [12, 14, 15].

Each patient should be treated in a method that is exactly suited for him or her. We have to watch the people who are experiencing issues and what their brains are like if they're not awake or if they're having convulsions. We should also assess the hearts of individuals who may have damaged their hearts, particularly if they are older or if they already have cardiac issues [13, 15]. If a person has inhaled smoke, we need to consider the possibility of toxic substances, such as cyanide, and treat them accordingly [5, 8].

Recently novel treatment techniques such as phototherapy, antioxidant approaches, and tailored ways to reduce oxidative stress and inflammation have been examined [9, 10]. These treatments are still primarily experimental, but they do suggest a growing interest in addressing pathways of harm beyond hypoxia alone. Early diagnosis and rapid treatment remain the most significant predictors of a good prognosis [9, 10, 15].

Delayed neurologic consequences:

Delayed neurological sequelae (DNS) are among the most severe effects of carbon monoxide poisoning and may arise days to weeks after apparent recovery. The most frequent signs are memory loss, cognitive impairment, mood changes, personality changes, gait abnormalities, and parkinsonian characteristics [7, 12, 15].

Chronic neuroinflammation, oxidative stress, demyelination, and delayed neuronal damage have been associated with the development of DNS. Abnormalities of the hippocampus, basal ganglia, and cerebral white matter are often seen in neuroimaging investigations [7, 10].

It is important to evaluate patients after first therapy since DNS may arise even with clinical recovery. Early identification of high-risk groups of patients may assist in decreasing long-term neurological damage [12, 15].

Prevention and Public Health:

Carbon monoxide poisoning is often prevented. Public health interventions should aim to raise knowledge of the hazards of exposure, encourage appropriate maintenance of fuel-burning equipment, and improve ventilation in enclosed spaces [4, 15]. The best preventative measure remains the installation of carbon monoxide monitors in homes and businesses. Occupational safety legislation and public education initiatives aimed at reducing the occurrence of accidental exposures and associated deaths are also critical [3, 4].

Emerging Research and Future Perspectives:

New research and views—progress in CO-related investigations—have enhanced insights into illness causes and diagnostic powers. Metabolomics analysis, novel biomarkers for oxidative stress, and state-of-the-art neuroimaging modalities might be potentially beneficial in early detection of severe instances of poisoning and more exact assessment of neurological prognosis [1, 7, 15]. The search is going underway for new therapy strategies targeting inflammation, poor mitochondrial activity, and destruction of brain cells [9,10].

In the future, modern breakthroughs in AI-supported diagnostics and precision medicine may be beneficial for clinical decision-making. Additional study on forensic markers and post-mortem diagnostics may also assist in a better identification of fatal instances [1, 13].

Conclusion:

Carbon monoxide poisoning is a frequent, avoidable cause of illness and death globally. It is not simply a result of hypoxia but also of other complex pathophysiologic processes such as oxidative stress, mitochondrial dysfunction, and inflammatory damage [5, 10, 15]. Difficulties in management are caused by the persistent difficulty with diagnosis, late neurological sequelae, and forensic concerns. Additional measures such as enhanced awareness and preventative procedures have a crucial role in the reduction of the incidence of carbon monoxide poisoning [14, 15].

References:

1. Ariza-Garcia, W. A., Pardo-Rodriguez, D., Taborda-Ocampo, G., Rosero-Moreano, M. & Cala, M. P. (2023). Metabolomic trends in deadly carbon monoxide poisoning: a forensic toxicology view. *Forensic Science International*, 377, 112643. *Forsciint* 2025 112643
<https://doi.org/10.1016/j.forsciint.2025>
2. Ruas, F., Mendonça, M. C., Real, F. C., Vieira, D. N., Teixeira, H. M. (2014). Carbon monoxide poisoning as a cause of death and differential diagnosis in the forensic practice: Retrospective research, 2000-2010. *Journal of Forensic and Legal Medicine*, 24, 1-6.
<https://doi.org/10.1016/j.jflm.2014.02.002>

3. Costa, M., Silva, B. S., Real, F. C., Teixeira, H. M. (2019). Epidemiology and forensics of carbon monoxide poisoning in Portugal: a three years investigation. *Forensic Science International*, 299, 1-5. 2019.03.016
4. Chen, Z., Liu, N., Tang, H., Gao, X., Zhang, Y., Kan, H. et al. (2022). Health impacts of exposure to sulfur dioxide, nitrogen dioxide, ozone and carbon monoxide from 1980 to 2019: a systematic review and meta-analysis. *Indoor Air* 32 (11).
<https://doi.org/10.1111/ina.13170>
5. Chenoweth, J. A., Albertson, T. E. & Greer, M. R (2021). Carbon monoxide poisoning. *Critical Care Clinics*, 37(3), 657-672.
<https://doi.org/10.1016/j.ccc.2021.03.010>
6. Ngorot Kembo, J. P., Wang, J., Luo, N., Gao, F., Yi, H., Zhao, S., et al. (2023). A review on catalytic oxidation of carbon monoxide over diverse catalysts with focus on hopcalite catalysts. *New Journal of Chemistry*, 47(44), 20222-20247.
<https://doi.org/10.1039/D3NJ03074A>
7. Li, W., Meng, J., Lei, J., Li, C., & Yue, W. (n. (2024). Carbon monoxide poisoning with hippocampi lesions on MRI: Case report and review of literature. *BMC Neurol.* 2024;24(1):159.
<https://doi.org/10.1186/s12883-024-03668-2>
8. Kao, L. W., Nañagas, K. A., & Penfound, S. J. (2022). Carbon monoxide poisoning. *Emerg Med Clin North Am.* 2022;40(2):283–312.
<https://doi.org/10.1016/j.emc.2022.01.005>
9. Zazzeron, L., Franco, W. and Anderson, R. (2024). Phototherapy; Carbon Monoxide Poisoning. *Nitric Oxide*, 146, 31-36.
<https://doi.org/10.1016/j.niox.2024.04.001>
10. Dent, M. R., Rose, J. J., Tejero, J., & Gladwin, M. T. (2024). Carbon monoxide poisoning: from micro-organisms to therapies *Annual Review of Medicine*, 75(1), 337-351.
<https://doi.org/10.1146/annurev-med-052422-020045>
11. Parmentier-Decrucq, E., & Mathieu, D. (2019). L'intoxication au monoxyde de carbone [Article en français]. *Revue du Praticien* 69(1):75-79. PMID: 30983293.
12. L. K. Weaver (2024). Carbon monoxide poisoning (Reprinted from the 2023 Hyperbaric Indications Manual, 15th edition). *Undersea & Hyperbaric Medicine*, 51(3), 253-276. PMID: 39348519.
13. Y., Ling, A., Ye, C., Xiong, X., Huang, H., & Wang, W. . (2026). Myocardial damage predicts death in elderly carbon monoxide poisoning: Development and internal validation of a simple predictive model. *Biomarkers*, 1-9.
<https://doi.org/10.1080/1354750X.2026.2681013>
14. Shih, R. D., Tomaszewski, C. A., Kaji, A., Diercks, D. B., Diercks, D. B., Anderson, J. D., et al. (2025). A important concern in the care of adult patients arriving to the emergency room with severe carbon-monoxide poisoning. *Annals of Emergency Medicine*, 85(4), e45-e59.
<https://doi.org/10.1016/j.annemergmed.2024.12.005>
15. Afzal, M., Agarwal, S., Elshaikh, R. H., Babker, A. M. A., Choudhary, R. K., Prabhakar, P. K., & et al. (2025). Carbon monoxide poisoning: pathogenesis, management, and future directions of therapy. *Diagnostics* 15(5), 581.
<https://doi.org/10.3390/diagnostics15050581>